

Tabelle 1: Hohe Anfälligkeit für das infektiöse Prionen Agens in MyD88<sup>-/-</sup> Mäusen nach peripherer und intracerebraler Inokulation.

	i.p. infection						i.c. infection					
	6 logLD <sub>50</sub>			3 logLD <sub>50</sub>			3x10 <sup>5</sup> LD <sub>50</sub>			3x10 <sup>2</sup> LD <sub>50</sub>		
Mouse genotype	At-tack rate	latency of disease in individual mice (days)	Mean ± SD (d)	At-tack rate	latency of disease in individual mice (days)	Mean ± SD (d)	At-tack rate	latency of disease in individual mice (days)	Mean ± SD (d)	At-tack rate	latency of disease in individual mice (days)	Mean ± SD (d)
MyD88 <sup>-/-</sup>	6/6	173, 181, 184	197±17	7/7	215, 225, 251	249±24	5/5	135, 135, 156	150±14	4/4	175, 177, 180	180±6
Wild-type	5/5	192, 206,	207±10	5/5	213, 222,	225±10	5/5	135, 135,	145±14	6/6	168, 168,	177±9
<i>Prnp</i> <sup>0/0</sup>	ND	-	-	0/5	> 370	> 370	ND	-	-	0/4	> 370	> 370