

Publikationen 2004

CRASH Trial Collaborators. Effect of intravenous corticosteroids on death within 14 days in 10008 adults with clinically significant head injury (MRC CRASH trial): randomised placebo-controlled trial. *Lancet* 2004;**364**:1321–8

Neff TA, **Stocker R**, Spahn DR. Fluid resuscitation in the intensive care unit. (Authors Reply) *N Engl J Med*. 2004;28:35:1905-8

Schmid A, Pugin J, Chevrolet JC, Marsch S, Ludwig S, **Stocker R**, Finnern H. Burden of illness imposed by severe sepsis in Switzerland. *Swiss Med Wkly*. 2004;134:97-102

Neff TA, **Stocker R**. Stable creatinine clearance using large-dose HES versus reduced GFR. *Kidney Int*. 2004;65:1111-2

Thomale UW, Griebenow M, Kroppenstedt SN, Unterberg AW, **Stover JF**. Small volume resuscitation with HyperHaes improves pericontusional perfusion and reduces lesion volume following controlled cortical impact injury in rats. *J Neurotrauma*. 2004; 21: 1737-46.

Unterberg AW, **Stover J**, Kress B, Kiening KL. Edema and brain trauma. *Neuroscience*. 2004;129(4):1021-9. Review.

Stover JF, Sakowitz OW, Unterberg AW. Neuronal activity and cortical perfusion determined by quantitative EEG analysis and laser doppler flowmetry are uncoupled in brain injured rats. *Acta Neurochir Suppl*. 2004;89:81-5.

Stover JF, Sakowitz OW, Kroppenstedt SN, Thomale UW, Kempinski OS, Flugge G, Unterberg AW. Differential effects of prolonged isoflurane anesthesia on plasma, extracellular, and CSF glutamate, neuronal activity, 125I-Mk801 NMDA receptor binding, and brain edema in traumatic brain-injured rats. *Acta Neurochir (Wien)*. 2004; 146: 819-30.

Hoover RC, Motta M, Davis J, Saatman KE, Fujimoto ST, Thompson HJ, **Stover JF**, Dichter MA, Twyman R, White HS, McIntosh TK. Differential effects of the anticonvulsant topiramate on neurobehavioral and histological outcomes following traumatic brain injury in rats. *J Neurotrauma*. 2004; 21: 501-12.