

[Features of cell death, mitochondrial activation and caspase dependence of rabbit anti-T-lymphocyte globulin signaling in lymphoblastic Jurkat cells are distinct from classical apoptosis signaling of CD95.](#)

Ziegler C, Finke J, Grüllich C.

Leuk Lymphoma. 2016;57(1):177-82. doi: 10.3109/10428194.2015.1044449.

Epub 2015 May 25.

PMID: 25927246

[Rabbit anti-T-lymphocyte globulin \(ATG\) persists with differential reactivity in patients' sera after full hematopoetic regeneration from allogeneic stem cell transplantation.](#)

Ziegler C, Finke J, Grüllich C.

Transpl Immunol. 2014 May;30(4):136-9. doi: 10.1016/j.trim.2014.03.005. Epub 2014 Apr 13.

PMID: 24727089

[Rabbit anti T-lymphocyte globulin induces apoptosis in peripheral blood mononuclear cell compartments and leukemia cells, while hematopoetic stem cells are apoptosis resistant.](#)

Grüllich C, Ziegler C, Finke J.

Biol Blood Marrow Transplant. 2009 Feb;15(2):173-82. doi:

10.1016/j.bbmt.2008.11.014.

PMID: 19167677

[Hepatitis B virus mutations associated with fulminant hepatitis induce apoptosis in primary Tupaia hepatocytes.](#)

Baumert TF, Yang C, Schürmann P, Köck J, Ziegler C, Grüllich C, Nassal M, Liang TJ, Blum HE, von Weizsäcker F.

Hepatology. 2005 Feb;41(2):247-56. doi: 10.1002/hep.20553.

PMID: 15660384